

19



Europäisches Patentamt
European Patent Office
Office européen des brevets

11

Publication number:

**0 337 354
A1**

12

EUROPEAN PATENT APPLICATION

21

Application number: 89106323.2

51

Int. Cl.⁴: **C11D 1/66 , C11D 3/37 ,
A61K 7/06**

22

Date of filing: 10.04.89

30

Priority: 12.04.88 JP 88291/88

43

Date of publication of application:
18.10.89 Bulletin 89/42

84

Designated Contracting States:
AT CH DE ES FR GB LI NL

71

Applicant: **Kao Corporation**
1-14-10, Nihonbashi Kayaba-cho
Chuo-ku Tokyo(JP)

72

Inventor: **Kamegai, Jun**
2-3-28, Baraki
Ichikawa-shi Chiba-ken(JP)
Inventor: **Hanazawa, Hideyuki**
3-20-1, Innai
Funabashi-shi Chiba-ken(JP)
Inventor: **Hirota, Hajime**
5-31-9, Megurohoncho Meguro-ku
Tokyo(JP)

74

Representative: **Wächtershäuser, Günter, Dr.**
Tal 29
D-8000 München 2(DE)

54

Low-irritation detergent composition.

57

A low-irritation detergent composition comprising (A) an alkyl saccharide-type surface active agent and (B) a cationic polymer. The composition is a low irritant to the skin and hair, produces excellent creamy foam, possesses a superior hair conditioning effect, and imparts an outstanding moisture feeling to the skin.

EP 0 337 354 A1

LOW-IRRITATION DETERGENT COMPOSITION

BACKGROUND OF THE INVENTION5 Field of the Invention:

This invention relates to a low-irritation detergent composition, and, more particularly, to a low-irritation detergent composition comprising an alkyl saccharide-type surface active agent and a cationic polymer. The composition is a low irritant to the skin and hair, produces excellent creamy foam, possesses a superior hair conditioning effect, and imparts an outstanding moisture feeling to the skin.

Description of the Background

15 Nonionic surface active agents are widely used in compositions for use in washing the skin and hair. Although nonionic surface active agents possess the advantage of being only slightly irritating, their foaming capability is not sufficient. Because of this, not much of them are formulated into detergent compositions requiring good foaming capability, such as hair shampoos, body shampoos, and the like. Besides the inferior foaming capability and relatively poor detergent capability, which are experienced when a nonionic surface active agent is used alone in a detergent composition, nonionic surface active agents have a problem of imparting a creaky, crinkle feeling to the hair and the skin.

Development of a stable detergent composition comprising a nonionic surface active agent, which is a low irritant to the skin and hair, possesses sufficient foaming and detergent capabilities, and is able to impart a superior feeling the skin, has been desired.

25 In view of this situation the present inventors have undertaken extensive studies to resolve the above-mentioned problem existing in nonionic surface active agents. As a result, the inventors have found that the combined use of an alkyl saccharide, which is one type of nonionic surface active agent, and a cationic polymer could greatly reduce irritation to the skin and hair, and produce abundant, high quality foam which impart a comfortable slippery feeling to the skin. Such a combined use of an alkyl saccharide and a cationic polymer also provided a detergent composition having an excellent hair conditioning effect and skin moisturizing effect.

SUMMARY OF THE INVENTION

35

Accordingly, an object of this invention is to provide a low-irritation detergent composition comprising the following components (A) and (B):

(A) at least one alkyl saccharide-type surface active agent represented by formula (I):

40 $R_1-O-(R_2O)_m-(G)_n$ (I)

wherein R_1 represents a linear or branched alkyl group of a C_8-18 carbon atom content, a linear or branched alkenyl group of a C_8-18 carbon atom content, or an alkylphenyl group of a C_8-18 carbon atom content, with the alkyl group being either linear or branched, R_2 represents an alkylene group of a C_2-4 carbon atom content, G represents a reduced sugar of a C_5-6 carbon atom content, m denotes a value of 0 to 10 and n denotes a value of 1 to 10, and

45 (B) at least one cationic polymer.

Other objects, features and advantages of the invention will hereinafter become more readily apparent from the following description.

50

DETAILED DESCRIPTION OF THE INVENTION AND PREFERRED EMBODIMENTS

As an alkyl saccharide-type surface active agent of formula (I), which is the (A) component of this

invention, those having an alkyl group for R_1 with 8 to 18, particularly 10 to 14 (decyl group, lauryl group, myristyl group, etc.), carbon atoms are preferable. The value of m in formula (I) is preferably from 0 to 3, with 0 being particularly preferable. The basic unit for the saccharide portion [G in formula (I)], which is the hydrophilic group of the alkyl saccharide-type surface active agent, is a reducing sugar having a C_5 - C_6 carbon atom content. Glucose, galactose, and fructose are named as examples of desirable reducing sugars. The degree of the polymerization (S) of saccharide, i.e., the value of n in formula (I), is 1 to 10. In particular, use of reducing sugars containing 80% or more of those having the degree of the polymerization (S) of 1 to 4 is desirable. The compounds of formula (I) having a lower degree of the polymerization (S), e.g. 1 to 1.4, are desirable. When the property of the compounds of formula (I) due to the group R_1 is taken into account, the value for the polymerization (S) of 1 to 1.4 is desirable for the R_1 group with C_8 - C_{11} , and (S) of 1.5 to 4.0 is desirable for the R_1 group with C_{12} - C_{14} . The mean values for (S) were determined by proton-NMR method.

Given as specific examples of these alkyl saccharidetype surface active agents are β -alkyl saccharide synthesized by the Koenigs-Knorr method such as octylglucoside, nonylglucoside, decylmaltoside, dodecylmaltoside, tridecylmaltoside, polyoxyethylene (2 E.O.)dodecylglucoside, and the like; alkyl saccharide produced from a reduced sugar such as glucose, galactose, maltose, or the like and a higher alcohol, polyoxyethylene alkylether glycol, or the like (USP 3,219,656, USP 3,839,318, and USP 4,223,129); and the like.


The (A) component is formulated into the detergent composition of this invention in an amount of 1 to 60% by weight. When the composition is a shampoo, an amount of 5 to 20% by weight is desirable. When it is a composition for use with the skin, an amount of 5 to 50% by weight is desirable.

Enumerated as examples of cationic polymers which can be used in this invention are: cationic cellulose derivatives, cationic starches, copolymers of a diallyl quaternary ammonium salt and an acryl amide, quaternarized polyvinylpyrrolidone derivatives, quaternarized vinylpyrrolidone vinylimidazol polymers (e.g. Luviscat, manufactured by BASF), polyglycol amine condensates, quaternarized collagen polypeptide, polyethylene imine, cationized silicon polymer [e.g. Amodimethicone, described in CTFA; cationic silicon polymers provided in a mixture with other components under the tradename of Dow Corning 929 (cationized emulsion), manufactured by Dow Chemical Co.], copolymers of adipic acid and dimethylaminohydroxypropyl diethylenetriamine (e.g. Cartaretine, manufactured by Sandoz Inc.), polyaminopolyamide (e.g. polymers described in French Patent Publication No. 2,252,840 and their crosslinked water-soluble polymers), cationic chitin derivatives, cationized guar gum (e.g. Jaguar C-B-S, Jaguar C-17, Jaguar C-16, etc. manufactured by Celanese Plastics and Specialties Company), quaternary ammonium salt polymers (e.g. Mirapol A-15, Mirapol AD-1, Mirapol AZ-1, etc., manufactured by Miranol Chemical Company Inc., U.S.A.; and polymers described in U.S.P. 2,261,002, U.S.P. 2,271,378, U.S.P. 2,273,780, U.S.P. 2,388,614, U.S.P. 2,454,547, U.S.P. 3,206,462, etc.), as well as cationic polymers selected from groups (a), (b), and (c) below.

(a) A polymer produced by reacting a polyaminopolyamide with epichlorohydrin at an epichlorohydrin mole ratio to the secondary amine group of the polyaminopolyamide of 0.5:1 to 1.8:1. This polyaminopolyamide is produced by the reaction of a polyalkylene-polyamine and a dicarboxylic acid selected from the members consisting of saturated fatty acid dicarboxylic acids having a C_3 - C_8 carbon atom content, and consisting of diglycol acid at a mole ratio of the polyalkylene-polyamine to the dicarboxylic acid of 0.8:1 to 1.4:1. Such a polymer is described in U.S.P. 3,227,615 and U.S.P. 2,961,347.

Specifically, this type of polymer is available from Hercules Corp. under the tradename of Hercoset 57. A 10% aqueous solution of this product has a viscosity of 30 cps. In particular, adipic acid epoxypolydiethylenetriamine copolymer is provided by Hercules Corp. under the tradename of PD170 or Delsette 101.

(b) A homopolymer or copolymer derived from an acrylic acid or a methacrylic acid having either of the following unit:



wherein R'_3 a hydrogen atom or an methyl group, A_1 represents a linear or branched alkyl group having a C_{1-6} carbon atom content or a hydroxylalkyl group having a C_{1-4} carbon atom content, R'_4 , R'_5 , and R'_6 , and which may be the same or different, represent an alkyl group having a C_{1-18} carbon atom content or a benzyl group, R'_1 and R'_2 individually represent a hydrogen atom or an alkyl group having a C_{1-6} carbon atom content, and X_1 represents a methosulfate anion or a halogen compound such as chloride or bromide.


Copolymers or mixtures of copolymers which can be used are those belonging to the group consisting of acrylamide, methacrylamide, diacetone-acrylamide, acrylamide or methacrylamide with a substituted lower alkyl group for the nitrogen atom, alkyl esters of acrylic or methacrylic acid, vinylpyrrolidone, and vinyl esters.

The following copolymers are given as specific examples. Products described in Cosmetic Ingredient Dictionary under the designation of QUATERNIUM 38, 37, 49, or 42; acrylamide/ β -methacryloyloxyethyl-trimethylammonium methosulfate copolymer provided by Hercules Corp. under the tradename of Reten 205, 210, 220, or 240; aminoethylacrylate phosphate/acrylate copolymer available from National Starch Co. under the tradename of Catrex, of which an 18% aqueous solution has a viscosity of 700 cps at 25°C ; and crosslinked cationic graft copolymers having a molecular weight of 10,000 to 1,000,000, preferably 15,000 to 500,000, and obtained copolymerizing (i) at least one type of monomer used for cosmetics, (ii) dimethylaminoethyl methacrylate, (iii) polyethylene glycol, and (iv) poly-unsaturated crosslinking agent. All of the above-mentioned copolymers are described in French Patent No. 2,189,434.


Crosslinking agents which can be used are selected from the group consisting of ethylene glycol dimethacrylate, diallyl phthalate, divinylbenzene, tetraallyl oxyethane, and polyallyl sucrose having 2 to 5 allyl groups per mole of sucrose.

The above-mentioned monomers used for cosmetics encompass a wide variety of monomers. Examples include vinyl esters of an acid having 2 to 18 carbon atoms, allyl- or methallyl esters of an acid having 2 to 18 carbon atoms, acrylate or methacrylate of a saturated alcohol having 1 to 18 carbon atoms, alkyl vinyl ether with the alkyl group having 2 to 18 carbon atoms, olefins having 4 to 18 carbon atoms, vinyl-type heterocyclic derivatives, dialkyl- or N,N-dialkylaminoalkyl maleate with the alkyl groups having 1 to 3 carbon atoms, and unsaturated acid anhydrides.

(c) Polymers comprising the units represented by the following formulae (I)' or (II)':



wherein A_2 is a group containing 2 amine groups, preferably piperidinyl groups, and Z represents the symbol B or B' , wherein B and B' may be the same or different and represent a linear or branched alkylene group which may be substituted with a hydroxyl group and may contain one or more oxygen atom, nitrogen atom, sulfur atom, 1 to 3 aromatic groups, and/or heterocyclic ring.




wherein A_2 has the same meaning as defined above, and Z_1 represents a symbol B_1 or B'_1 , at least one Z_1 being the B'_1 , wherein B_1 represents a linear or branched alkylene or hydroxy alkylene group and B'_1 represents a linear or branched alkylene group which has at least one nitrogen atom bonded in the chain and may have one or more hydroxyl group substituents. The nitrogen group has a substituent alkyl group which may have an oxygen atom(s) in the chain and a hydroxyl group substituent(s).

Included also in this group of polymers are cationic polymers selected from quaternary ammonium salts or oxidized products of the above polymers (I)' or (II)'.


Various cationic polymers can be enumerated as mentioned above. Preferable cationic polymers, which

can be used as the (B) component of this invention, however, are cationic cellulose derivatives, cationic starches, copolymers of a diallyl quaternary ammonium salt and an acryl amide, quaternarized polyvinylpyrrolidone derivatives, and polyglycol amine condensates.

As a cationic cellulose derivative, a compound which satisfies the following formula (II) is desirable:




wherein A indicates a residual group of anhydroglucose units, 1 is an integer of 50 to 20,000, and each R_3 indicates a substitution group shown in the following formula (III):



wherein R' and R'' represent alkylene groups having 2 or 3 carbon atoms, o is an integer of 0 to 10, o' is an integer from 0 to 3, p is an integer of 0 to 10, R'' represents an alkylene group or hydroxyalkylene group having 1 to 3 carbon atoms, R_4 , R_5 , and R_6 may be the same or different, and represent an alkyl, aryl, or aralkyl group having not more than 10 carbon atoms, or may form a heterocyclic ring with a nitrogen atom in the formula, and X designates an anion (chlorine, bromine, iodine, sulfuric acid, sulfonic acid, methyl-sulfuric acid, phosphoric acid, nitric acid, and the like).

The degree of cation substitution of the cationic cellulose, that is, the average value of o' for each anhydroglucose unit is preferably between 0.01 to 1, with the range of 0.02 to 0.5 being more preferable. Also, the total of $o + p$ averages between 1 and 3. A substitution value of smaller than 0.01 is unsatisfactory, while if this value is greater than 1 there is no particular problem, but, from the aspect of reaction yield, a value 1 or smaller is more desirable. The molecular weight of the cationic cellulose used here is in the range of about 100,000 to 3,000,000.



A desirable cationic starch for use in the present invention should satisfy the following formula (IV):



wherein A' represents a starch residual group, R_7 represents an alkylene group or hydroxyalkylene group, R_8 , R_9 , and R_{10} may be the same or different, and represent an alkyl, aryl, or aralkyl group having less than 10 carbon atoms, or may form a heterocyclic ring together with a nitrogen atom in the formula, X' represents an anion (chlorine, bromine, iodine, sulfuric acid, sulfonic acid, methyl-sulfuric acid, phosphoric acid, nitric acid, and the like.), and 1' is a positive integer.

The degree of cation substitution of the cationic starch, i.e., the number of cation groups for each anhydrous glucose group, is desirably 0.01 to 1, with the more desirable range being 0.02 to 0.5. A substitution value of less than 0.01 is unsatisfactory, while if this value is greater than 1 there is no particular problem. However, from the aspect of reaction yield, a value 1 or smaller is more desirable.

A desirable copolymer of a cationic diallyl quaternary ammonium salt and acryl amide for use in the present invention is that which satisfies the following formulae (V) or (VI):



25


30

wherein R_{11} and R_{12} , which may be the same or different, represent a hydrogen, an alkyl group having 1 to 18 carbon atoms, a phenyl group, an aryl group, a hydroxyalkyl group, an amidoalkyl group, a cyanoalkyl group, an alkoxyalkyl group, or a carboalkoxyalkyl group, R_{13} , R_{14} , R_{15} , and R_{16} , which may be the same or different, represent a hydrogen, a lower alkyl group having 1 to 3 carbon atoms, or a phenyl group, X^{\ominus} represents an anion (chlorine, bromine, iodine, sulfuric acid, sulfonic acid, methyl-sulfuric acid, phosphoric acid, nitric acid, or the like.), l_1 is an integer of 1 to 50, m_1 is an integer of 1 to 50, and n_1 is an integer of 150 to 8,000.

35

The molecular weights of copolymers of diallyl quaternary ammonium salt and acryl amide may be in the range from about 30,000 to 2,000,000, but the range from 100,000 to 1,000,000 is particularly preferable.

A desirable quaternary polyvinylpyrrolidone derivative which can be used in the present invention is represented by the following formula (VII):



50


55

wherein R_{17} represents a hydrogen atom or an alkyl group having 1 to 3 carbon atoms, R_{18} , R_{19} , and R_{20} , which may be the same or different, represents a hydrogen atom, an alkyl group, a hydroxyalkyl group, an amidoalkyl group, a cyanoalkyl group, an alkoxyalkyl group, or a carboalkoxyalkyl group, with the alkyl groups having 1 to 4 carbon atoms, Y represents an oxygen atom or an NH group in an amide group, X^{\ominus} designates an anion (chlorine, bromine, iodine, sulfuric acid, sulfonic acid, alkylsulfuric acid having 1 to 4 carbon atoms, phosphoric acid, nitric acid, and the like.), q is an integer from 1 to 10, and m_2 and n_2 represent integers satisfying the equation $m_2 + n_2 = 20-8000$.

The molecular weights of the quaternary polyvinyl pyrrolidone derivative may be in the range from about 10,000 to 2,000,000, but the range from 50,000 to 1,500,000 is particularly desirable.

The content of the cationic nitrogen derived from the cationic polymer contained in the above-mentioned vinyl polymer is in the range from 0.004% to 0.2% by weight, with the most desirable range being from 0.01% to 0.15% by weight. Below 0.004% the effect is inadequate; above 0.2% the performance is good, but the vinyl polymer becomes colored and the high cost is disadvantageous.

5 Given as a desirable polyglycol-polyamine condensate a compound represented by the following formula (VIII):



wherein R₂₂, R₂₄, R₂₅, and R₂₇ represent hydroxyalkylene groups having 2 to 4 carbon atoms, R₂₃ and R₂₆ represent alkylene groups having 2 or 3 carbon atoms, x and y represent integers of 10 to 20, m₃ represents an integer of 2 to 4, n₃ represents an integer of 2 to 6, r represents an integer of 1 to 50, and R₂₁ represents a linear or branched alkyl group having 6 to 20 carbon atoms.

One or more of the above-mentioned cationic polymers are used as the (B) component and are formulated into the detergent composition of this invention in an amount of 0.01 to 5% by weight. When the composition is a shampoo the amount of 0.1 to 1.5% by weight is desirable. When it is a composition for use with the skin the amount of 0.01 to 1% by weight is desirable.

It is desirable to adjust the pH of the composition of this invention to 2-10, preferably to 4-8, using an acidic or alkaline pH adjusting agent which is conventionally used in a detergent composition.

Besides the above-mentioned essential components, other components which are generally used in detergent compositions may be optionally formulated into the detergent composition of the present invention inasmuch as the effect of the present invention is not affected. Such components include, for example, humectants such as propylene glycol, glycerin, sorbitol, and the like; viscosity adjusting agents such as carboxyvinyl polymer, methyl cellulose, hydroxyethyl cellulose, polyoxyethyleneglycol distearate, ethanol, and the like; pearling agents; perfumes; pigments; ultraviolet ray absorbers; antioxidants; biocidal agents (Trichlosan, Trichlorocarban, etc.); antiphlogistic agents (potassium glycol phosphate, tocopherol acetate, etc.); anti-dandruff agents (zinc pyrithione, Octopirox, etc.); antiseptics (methyl paraben, butyl paraben, etc.); and the like.

The detergent composition of this invention can take any conventionally employed preparation forms. The desirable proportion of the components (A) plus (B) in the total amount of the surface active agents in the composition is 30% by weight or greater, when the composition is a solid form, and 10% by weight or greater, when the composition is a liquid form.

Other features of the invention will become apparent in the course of the following description of the exemplary embodiments which are given for illustration of the invention and are not intended to be limiting thereof.

40 EXAMPLES

45 Example 1

Shampoos having formulations listed in Table 1 were prepared. Foaming capability and feeling upon touch were evaluated for each shampoo. The results are presented in Table 1.

50 <Evaluation Method>

One (1) gm of a shampoo was applied to tresses of hair taken from healthy Japanese women, 15 cm long, weighing 20 gm. The shampoo was lathered for 1 minute, and the foaming conditions were evaluated by 5 expert panelists. Then, the tress was rinsed with tap water for 30 seconds at 30 °C, and dried first with a towel, and then with a dryer. The feeling upon touch of the dried hair was evaluated by the same panelists.

<Evaluation standard>

Foaming capability:

5

AAA: Lather was abundant

BBB: Lather was normal

CCC: Lather was insufficient

10 Quality of Lather:

AAA: Lather was creamy

BBB: Lather was rough

15 Feeling to touch:

AAA: Hair felt soft and not creaky

BBB: Hair was not slippery and felt creaky

BBB: Hair was not slippery and felt creaky

20

TABLE 1

25	(components: % by weight)						
		Comparative Product		Inventive Product			
		1	2	1	2	3	4
30	C ₁₀ -O-(G) _{1.5} * 1	20	20	20	20	20	20
	C ₁₂ -O-(G) _{2.0} * 2			0.5	0.5		
	Polymer JR-400 * 3					0.5	
	Cationized starch						0.5
	Merquat 550 * 4						
35	Gafquat 755 * 5						
	Purified water	80	80	79.5	79.5	79.5	79.5
	Evaluation of lather						
	Foaming capability	BBB	BBB	AAA	AAA	AAA	AAA
	Quality of lather	CCC	CCC	AAA	AAA	AAA	AAA
40	Feeling upon touch	CCC	CCC	AAA	AAA	AAA	AAA

* 1 C₁₀: decyl group; G: glucose

* 2 C₁₂: dodecyl group; G: glucose

* 3 A cationized cellulose, manufactured by Union Carbide Corp.

45 * 4 A copolymer of N,N-dimethyl-3,5-methylenepiperidinium chloride and acrylamide; manufactured by Merck Co.

* 5 A triethanolamine salt of ethyl sulfate quarternirized product of vinylpyrrolidone/dimethylaminoethyl acrylate manufactured by Guf Chemical Co.

50

Examples 2

Five percent (5%) aqueous solutions of the detergent compositions of this invention prepared as in Example 1 were subjected to a closed patch test for 24 hours by 5 volunteered panellists. None of the solutions were felt to irritate the skin. Thus, the composition of the invention was proven to be a low irritant.

55

Example 3

Paste face cleansing foam:

<Formulation>

5

	Components	% by weight
10	(1) Alkyl saccharide [C ₁₂ -O-(G) _{2.0}] *	45
	(2) Cationized cellulose (Polymer JR-400, manufactured by Union Carbide Corp.)	1
	(3) Ethylene glycol distearate	3
	(4) Polyethylene glycol 400	0.5
	(5) Methyl paraben	0.2
15	(6) Perfume	0.2
	(7) Purified water	Balance

* C₁₂: dodecyl group; G: glucose

20

<Preparation>

To water heated to 70 °C were added components (1) to (5) under stirring to obtain a transparent solution. After cooling this solution to 40 °C, component (6) was added and the mixture was cooled to the room temperature.

The pasty face cleansing foam containing an alkyl saccharide-type surface active agent and a cationized cellulose thus prepared produced abundant lather, was able to be readily rinsed away, and provided a comfortable moistening feeling after cleansing. In addition, the face cleansing foam had a superior low-temperature stability.

30

Example 4

35 Shampoo composition:

<Formulation>

40

	Components	% by weight
	(1) Alkyl saccharide [C ₁₀ -O-(G) _{1.5}] * 1	20
	(2) Merquat 550 * 2	0.3
45	(3) Polyethylene glycol distearate * 3	1
	(4) Methyl paraben	0.2
	(5) Blue #1	Small amount
	(6) Perfume	0.2
50	(7) Purified water	Balance

* 1 C₁₀: decyl group; G: glucose

* 2 A copolymer of N,N-dimethyl-3,5-methylenepiperidinium chloride and acrylamide; manufactured by Merck Co.

* 3 Emanone 3299 manufactured by Kao Corp.

55

<preparation>

To purified water heated to 60°C were added components (1) to (5) under stirring to obtain a transparent solution. After cooling this solution to 40°C, component (6) was added and the mixture was cooled to room temperature.

The shampoo composition containing an alkyl saccharide-type surface active agent and a cationized polymer thus prepared produced abundant lather, was able to be readily rinsed away, and provided a comfortable moistening feeling to the hair after shampooing. In addition, the shampoo composition had a superior low-temperature stability.

Example 5

Shampoo composition

<Formulation>

Components	% by weight
(1) Alkyl saccharide [C ₁₀ -O-(G) _{1.4}] * 1	10
(2) Gafquat 755 * 2	0.4
(3) Sodium polyoxyethylene(2)lauryl sulfate	5
(4) Coconut oil diethanolamide	1
(5) Sodium chloride	1
(6) Perfume	0.2
(7) Purified water	Balance

* 1 C₁₀: decyl group; G: glucose

* 2 A ethyl sulfate quarternirized triethanolamine of vinylpyrrolidone/dimethylaminoethyl acrylate manufactured by Gaf Chemical Co.

To purified water heated to 60°C were added components (1) to (5) under stirring to obtain a transparent solution. After cooling this solution to 40°C, component (6) was added and the mixture was cooled to room temperature.

The shampoo composition containing alkyl saccharide-type surface active agent and a cationized polymer thus prepared produced abundant lather, was readily rinsed away, and provided a comfortable moistening feeling to the hair after shampooing. In addition, the shampoo composition had a superior low-temperature stability.

Example 6

Anti-dandruff shampoo

<Formulation>

Components	% by weight
(1) Alkyl saccharide [C ₁₂ -O-(G) _{1.7}] *	15
(2) Cationized cellulose (Polymer JR-400, manufactured by Union Carbide Corp.)	0.2
(3) Tomicide Z-50 (Zinc pyrithion, manufactured by Yoshitomi Pharmaceutical Ind.)	1
(4) Methyl paraben	0.1
(5) Perfume	0.2
(6) Purified water	Balance

* C₁₂: lauryl group; G: glucose

To purified water heated to 60° C were added components (1) to (4) under stirring to obtain a transparent solution. After cooling this solution to 40° C, component (5) was added and the mixture was cooled to room temperature.

The anti-dandruff shampoo containing alkyl saccharidetype surface active agent and a cationized polymer thus prepared produced abundant lather, was readily rinsed away, and provided a comfortable moistening feeling to the hair after shampooing. In addition, the shampoo composition had a superior low-temperature stability.

Example 7

Liquid face cleansing form

<Formulation>

Components	% by weight
(1) Alkyl saccharide [C ₁₂ -O-(G) _{1.7}] *	35
(2) Cationized cellulose (Polymer JR-400, manufactured by Union Carbide Corp.)	1
(3) Ethylene glycol distearate	3
(4) Triethanolamine laurate	5
(5) Methyl paraben	0.2
(6) Ethanol	4
(7) Perfume	0.2
(8) Purified water	Balance

* C₁₂: lauryl group; G: glucose

To purified water heated to 70° C were added components (1) to (5) under stirring to obtain a transparent solution. After cooling this solution to 50° C, component (6) was added and the mixture was cooled to room temperature. Component (7) was then added to the mixture.

The liquid face cleansing form containing alkyl saccharide-type surface active agent and a cationized polymer thus prepared produced abundant lather, was readily rinsed away, and provided a comfortable moistening feeling to the skin after cleansing. In addition, the liquid face cleansing form had a superior low-temperature stability.

Example 8

Body shampoo

<Formulation>

Components	% by weight
(1) Alkyl saccharide [C ₁₂ -O-(G) _{1.7}] * 1	30
(2) Merquat 550 * 2	0.1
(3) Ethylene glycol distearate	3
(4) Hydroxyethyl cellulose	0.2
(5) Irgasan DP-300 (Triclosan, manufactured by Ciba Geigy)	0.5
(6) Ethanol	5
(7) Methyl paraben	0.2
(8) Perfume	0.2
(9) Purified water	Balance

* 1 C₁₂: lauryl group; G: glucose

* 2 A copolymer of N,N-dimethyl-3,5-methylenepiperidinium chloride and acrylamide; manufactured by Merck Co.

To purified water heated to 70 °C were added components (1) to (7) under stirring to obtain a transparent solution. After cooling this solution to 40 °C, component (8) was added and the mixture was cooled to room temperature.

The body shampoo containing alkyl saccharide-type surface active agent and a cationized polymer thus prepared produced abundant lather, was readily rinsed away, and provided a comfortable moistening feeling to the skin after shampooing. In addition, the body shampoo had a superior low-temperature stability.

Example 9

Shampoo composition

<Formulation>

Components	% by weight
(1) Alkyl saccharide [C ₈ -O-(G) _{1.3}] * 1	20
(2) Jaguar C-13-S * 2	0.5
(3) Octopirox	0.5
(4) Disodium ethylenediamine tetraacetate	0.4
(5) Perfume	0.6
(6) Ethanol	5
(7) Purified water	Balance

* 1 C₈: octyl group; G: glucose

* 2 Quaternarized guar gum, manufactured by Celanese Co.

To purified water heated to 60 °C were added components (1) to (4) under stirring. Component (6) was added to the solution which was cooled to 55 °C. After cooling this solution to 40 °C, component (5) was added and the mixture was cooled to room temperature.

The shampoo composition containing alkyl saccharide-type surface active agent and a cationized polymer thus prepared produced abundant lather, was readily rinsed away, and provided a comfortable moistening feeling to the hair after shampooing.

Example 10

Shampoo composition

5

<Formulation>

10

Components	% by weight
(1) Alkyl saccharide $[C_{12}-O-(CH_2CH_2O)_2-G] \cdot 1$	18
(2) Jaguar C-13 * 2	0.5
(3) Laurylamineoxide	2.0
(4) Perfume	0.5
(5) Purified water	Balance

15

* 1 C_{12} : lauryl group; G: glucose

* 2 Cationized guar gum manufactured by Celanese Co.

20

To purified water heated to 70 °C were added components (1) to (3) under stirring. Component (4) was added to the solution which was cooled to 50 °C. This solution was cooled to room temperature to obtain the target shampoo composition.


25 The shampoo composition containing alkyl saccharide-type surface active agent and a cationized polymer thus prepared produced abundant lather, was readily rinsed away, and provided a comfortable moistening feeling to the hair after shampooing.

Obviously, numerous modifications and variations of the present invention are possible in light of the above teachings. It is therefore to be understood that within the scope of the appended claims, the invention 30 may be practiced otherwise than as specifically described herein.

Claims

35 1. A low-irritation detergent composition comprising the following components (A) and (B):

(A) at least one alkyl saccharide-type surface active agent represented by formula (I):



wherein R_1 represents a linear or branched alkyl group of a C_8-18 carbon atom content, a linear or branched alkenyl group of a C_8-18 carbon atom content, or an alkylphenyl group of a C_8-18 carbon atom content, with the alkyl group being either linear or branched, R_2 represents an alkylene group of a C_2-4 carbon atom content, G represents a reduced sugar of a C_5-6 carbon atom content, m denotes a value of 0 to 10 and n denotes a value of 1 to 10, and

(B) at least one cationic polymer.

45 2. A low-irritation detergent composition according to Claim 1, wherein said cationic polymer is at least one member selected from the group consisting of cationic cellulose derivatives, cationic starches, copolymers of a diallyl quaternary ammonium salt and an acryl amide, quaternarized polyvinylpyrrolidone derivatives, and polyglycol amine condensates.

50

55



DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
Y	GB-A-2 050 166 (L'OREAL) * example 16; claims 1,10-19 * ---	1,2	C 11 D 1/66 C 11 D 3/37 A 61 K 7/06
Y	US-A-4 048 301 (C. PAPANTONIOU) * abstract; column 1, lines 56-60; column 2, lines 4-9 ; claims * ---	1,2	
A	US-A-4 678 595 (A. H. MALIK) * abstract; column 3, lines 24-29; claims 1-3 * ---	1,2	
A	GB-A-2 128 627 (L'OREAL) * abstract; claims * ---	1,2	
A	US-A-4 493 773 (COOK et al.) * claims * -----	1,2	
			TECHNICAL FIELDS SEARCHED (Int. Cl.4)
			C 11 D A 61 K
The present search report has been drawn up for all claims			
Place of search BERLIN		Date of completion of the search 07-07-1989	Examiner PELLI-WABLAT B
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ----- & : member of the same patent family, corresponding document	